



Division of Infectious Diseases

400 Community Drive
Manhasset, NY 11030
516- 757-7934



Learn More About mpox Vaccine

Mpox is a virus that can cause a painful rash. It spreads mostly through prolonged skin-to-skin contact, especially sexual contact. It can also be spread by touching items that someone with mpox has touched.



If you think you might be at risk for getting mpox, contact us below about getting vaccinated.



Do not share eating utensils, cups, bowls, personal items, bed linens, towels, and clothes with people who have a rash that looks like mpox.



Avoid skin-to-skin and sexual contact with people who have a rash.



Wash your hands often with soap and water.



JYNNEOS is a 2-dose mpox vaccine

Vaccination is an important tool in preventing the spread of painful lesions due to mpox.

- Applied on the forearm and other injection area's are available.
- The second dose is given 4 weeks (28 days) after the first dose.
- An immune response is seen right after the first dose. Full protection is seen two weeks after the second dose.

The **most common side effects** are pain, redness, and itching at the injection site. Experiencing fever, headache, tiredness, nausea, chills, and muscle aches are common signs that your immune system is responding and working on building immunity! Some people have not exhibited any side effects.



Side Effects



- ▶ Scan here to read more information.
- ▶ Contact the Northwell Health mpox resource line
📞 833-315-1748
📍 24/7 hotline

**Schedule your vaccination
appointment with us today!**

📞 516- 757-7934

